

FTD-ID(RS)T-0792-88

# AD-A223 114


# FOREIGN TECHNOLOGY DIVISION



SPECIAL FEATURES OF THE STRONG INTERACTION OF A PLASMA WITH A MAGNETIC FIELD AS IT EXPANDS RADIALLY IN A DISK CHANNEL

by

A. Yu. Kerbis, V.S. Sokolov, et al.



Approved for public release; Distribution unlimited.

MICROFORM CLASSIFICATION NOTICE/TARGET (For Unclassified Material)

(TL) BE COMPLETED BY RECORDS CUSTODIAN OR CAMERA OPERATOR/DOCUMENT PREPARER IN GOVERNMENT OR CONTRACT MICROFORM SERVICE CENTER)


# CLASSIFICATION NOTICE THE HIGHEST CONTAINED ON THIS MICROFORM IS:

# **UNCLASSIFIED**

IF CHECKED THE FOLLOWING ACCESS WARNING A	NOTICES APPLY
 FOR OFFICIAL USE ONLY (AFR 12-30)	
PRIVACY ACT OF 1974 (AFR 12-35)	
SOURCE SELECTION SENSITIVE (AFR 70-15)	
OTHER (Specify)	(For Unclassified Material)

**AFSC Form 201, AUG 87** 

# DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

# **HUMAN TRANSLATION**

FTD-ID(RS)T-0792-88

25 October 1988

MICROFICHE NR: FTD-88-C-000888

SPECIAL FEATURES OF THE STRONG INTERACTION OF A PLASMA WITH A MAGNETIC FIELD AS IT EXPANDS RADIALLY IN A DISK CHANNEL

By: A. Yu. Kerkis, V.S. Sokolov, et al.

English pages: 6

Source: Novosibirsk Aerofizicheskiye Issledovaniya,

pp. 71-75

Country of origin: USSR

Translated by: Charles T. Ostertag, Jr.

Requester: FTD/TQTD

Approved for public release; Distribution unlimited.

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION FOREIGN TECHNOLOGY DIVISION WPAFB, CHIO.

FTD-ID(RS)T-0792-88

Date 25 October

1988

U. S. BOARD ON GEOGRAPHIC NAMES TRANSLITERATION SYSTEM

Blo	ock	Italic	Transliteration	Block	Italic	Transliteration
A	8	A 4	Α, α	Рр	P	R, r
Б	Q	5 6	В, Ъ	Cc	Cc	S, s
8	•	8 •	V, v	Ττ	7 m	T, t
Γ	Γ	<i>r</i> •	G, g	Уу	yy	U, u
Д	Д	4	D, d	Фф	• •	F, f
Ε	e	E .	Ye, ye; E, 🤳	Х×	Xx	Kh, kh
ж	m	<i>X</i> ×	ih, zh	Цц	4 .	Ts, ts
3	з.	3 ,	Z, z	4 4	4 4	Ch, ch
И	н	H W	I, i	Шш		Sh, sh
Я	й		Y, y	Щщ	Щщ	Shch, shch
н	H	KK	K, k	ъъ	3 .	H
Л	Л	A A	L, 1	ы н	M u	Y, y
M	m	M M	M, m	<b>b b</b>	<b>b</b> •	•
H	н	H M	N, n	Э э	<b>3</b> ,	E, e
C	0	0 •	0, 0	10 to	D .	Yu, yu
П	п	/7 A	P, p	Яя	Я	Ya, ya

\*ye initially, after vowels, and after b, b; e elsewhere. When written as & in Russian, transliterate as ye or &.

# RUSSIAN AND ENGLISH TRIGONOMETRIC FUNCTIONS

Russian	English	Russian	English	Russian	English
sin	sin	sh	sinh	arc sh	sinh <sup>-1</sup>
cos	COS	ch	cosh	arc ch	cosh <sup>-1</sup>
tg	tan	th	tanh	arc th	tanh 1
ctg	cot	cth	coth	arc cth	coth <sup>-1</sup>
sec	sec	sch	sech	arc sch	sech <sup>-1</sup>
COSC	CSC	csch	csch	arc cach	csch <sup>-1</sup>

Russian	English	
rot	cur1	
1g	log	

# GRAPHICS DISCLAIMER

All figures, graphics, tubles, equations, etc. merged into this translation were extracted from the best quality copy available.



# SPECIAL PEATURES OF THE STRONG INTERACTION OF A PLASMA WITH A MAGNETIC FIELD AS IT EXPANDS RADIALLY IN A DISK CHANNEL

Acces	sion for		
NTIS	GRA&I	12	
DTIC	TAB	ð	
Unani	ounced		
Just1	fication		
<del>`</del>			
Бу			
Distribution/			
 5\f	il abality C	ode <b>s</b>	
	azat md.	/or	
	1 2 Jac 1al		
ı	1		
. A	97		
H-/	مل		
	1		

Kerkis, A.Yu., Sokolov, V.S., Trynkine, N.A., Fonichev, V.P.,

The investigation concerns the process of expansion of a plasma in a disk channel when a transverse magnetic field is present. The layout of the experimental unit is shown in Figure 2.19. 7 The plasma source is a coaxial electric discharger, on which a capacitor bank of the IMH-5-150 type is discharged. The initial eir pressure comprises - > + × P 5 0.7-1.0 mm Hg. The plasma which forms as a result of the electric discharge moves along a straight tube, which is a continua. For of the discharger, at a velocity or up to 20 km/s. This tube, which was a length of 30 cm and diameter of 5 cm, terminates in a wage charmant with a width of 2 cm. The process develops in such a manner that the plasma, following impact on the wnd of the straight tube, is expanded in the disk channel. The leading edge of the moving plasma, already at a radius of 60 mm, is a shock wave, traveling over a quiescent gas at a velocity of up to 10 km/s (without a magnetic field). The results of the measurement of pressure along the length of the disk channel at different moments of time when the magnetic field is absent are given in Figure 2.20. Also given are the estimates and the indirect measurements of distribution of electrical conductivity over the radius at different moments of time. These measurements in combination with

data on pressure showed that the change in the parameters of the gas beyond the cylindrical shock wave take place primarily as a result of its adiabatic expansion.

In this case the presence of a transverse magnetic field, when its strength exceeds a certain critical sagnitude, leads to a qualitative change in the nature of expansion of the gas. Direct measurements of the current density in the plasma, changes in the strength of the magnetic and electric fields, pressure (Figure 2.21), and also high-speed photography with the help of an SFRL (in the mode of photosweep and time magnifier), lead to the detection of a clearly expressed current layer (Figure 2.22), appearing in the gas beyond the direct shock wave and leading, as a result of its intensive stagnation in the magnetic field, to the formation of a reflected shock wave. In this case the measured current density reaches a magnitude of 2000  $A/cm^2$ , which at a field strength of  $\sim 4000$  Oe under the conditions of the experiment leads practically to the complete standstill of the gas in this zone. Estimates show that the magnitude of electrical conductivity inside the current layer reaches values of 120-150 who/cm (without a magnetic field it is equal to 5-10 mho/cm).




Figure 2.19.

Key: (1) Pressure sensors; (2) Current sensors; (3) Magnetic probes;
(4) SFR.




Figure 2.20.




Figure 2.21.

Rey: (1) µs.




Figure 2.22. Frame-by-frame photography of the process. Exposure time for one frame 1.33 µs. Sequence of frames: from top to bottom, from left to right.

During the course of the experiment attention was turned to the stability of the current layer. It was revealed that it had exceptional stability as a geometric formation. In a number of cases on the external boundary of the layer a picture developed which was characteristic for Rayleigh-Taylor instability, which, however, disappeared quite rapidly, leading to the complete restoration of the structure of the current layer. The process of expansion of the conducting gas in a transverse magnetic field, which is being investigated at the present time, and which is accompanied by the formation of a clearly expressed current layer, coincides in its nature with the analogous process which was described in [1], and thereby confirms experimentally the conclusions of this work, obtained on the basis of a numerical solution of equations of magnetic hydrodynamics.

### BIBLIOGRAPHY

- 1. A.W. Tikhonov, A.A. Samerskiy, L.A. Zaklyaz'minukiy,
- 7.M. Volomevich, L.M. Degtyarev, S.P. Kurdyumov,
- Yu.P. Popov, V.S. Sokolov, A.P. Favorskiy, "Monlinear effect of the formation of a self-austaining, high-temperature conducting layer of gas in nonsteady-state processes of magnetic hydrodynamics," DAN [Reports of the Academy of Sciences], No 4, v 173, 1967.

# DISTRIBUTION LIST

# DISTRIBUTION DIRECT TO RECIPIENT

CRCANTIZATION	MICROFICHE
A205 DYAHIC	1
A210 DAAC	1
CS09 BALLISTIC RES LAB	
C510 RET LABS/AVEADCOM	1
C513 ARRADOOM	1
C535 AVRADOOM/TSAROOM	1
C539 TRASANA	1
C591 FSTC	4
C619 MIA REDSTONE	1
DOOR MISC	1
E053 HQ USAF/INET	1
E404 AEDC/DOF	ī
E408 AFWL	1
E410 AD/IND	1
E429 SD/IND	1
POOS DOE/ISA/DDI	1 2
POSO CIA/OCR/ADD/SD	
AFIT/LDE	1
FID	_
<b>∝</b> v	1
MIA/PHS	1
LLYL/CODE L-389	1
NASA/NST-44	1
NSA/T513/TDL	2
ASD/FTD/TQLA	1
FSL/NIX-3	1

# **MICROFORM CERTIFICATE OF AUTHENTICITY**

### START

THIS IS TO CERTIFY THAT THE MICROGRAPHIC DMAGES AFPEARING ON THIS MICKOPORM ARE ACCURATE REPPODUCTIONS
OF THE RECORDS CREATED OR MAINTAINED BY: THE FOREIGN TECHNOLOGY DIVISION,
WRIGHT-PATTERSON AFB, OHIO

### PREVIOUS DESIGNATIONS

AND WERE MICROFILMED IN THE COURSE OF BUSINESS PURSUANT TO ESTABLISHED ROUTINE POLICY FOR SYSTEMS
UTILIZATION AND/OR FOR THE MAINTENANCE AND PRESERVATION OF SUCH RECORDS THROUGH THE STORAGE OF SUCH
MICROFORMS IN PROTECTED LOCATIONS.

IT IS FURTHER CERTIFIED THAT THE PHOTOGRAPHIC PROCESSES USED FOR MICROFILMING OF THE ABOVE RECORDS WERE ACCOMPLISHED IN A MANNER AND ON MICROFILM THICH MEETS THE RECOMMEMDED REQUIREMENTS OF THE NATIONAL BUREAU OF STANDARDS FOR MICROGRAPHIC REPRODUCTIONS.

DATE MICROFILMED 21 Nov 88	CAMERA OPERATOR	
PLAL NUMBER	RECORDS MANAGEMENT DIVISION	
(3486) (2609) (1062)	FTD/DA	
FIL DIMENSEATION	241	
PTD/DARM, WPAFB, OHIO	REDUCTION RATIO	
AUTHORITY AFR 12-40 AND MICROFORM SYSTEM NUMBER		
APSC 1R-72		

AFSC Form 96, AUG 87